

Higher Curvature Corrections to Primordial Fluctuations in Slow-roll Inflation

Masaki Satoh* and Jiro Soda†

**†Department of Physics, Kyoto University, Kyoto 606-8501, Japan*

(Dated: June 27, 2008)

We study higher curvature corrections to the scalar spectral index, the tensor spectral index, the tensor-to-scalar ratio, and the polarization of gravitational waves. We find that the higher curvature corrections can not be negligible in the dynamics of the scalar field, although they are energetically negligible. Indeed, it turns out that the tensor-to-scalar ratio could be enhanced and the tensor spectral index could be blue due to the Gauss-Bonnet term. We estimate the degree of circular polarization of gravitational waves generated during the slow-roll inflation. We argue that the circular polarization can be observable with the help both of the Gauss-Bonnet and parity violating terms. We also present several examples to reveal observational implications of higher curvature corrections for chaotic inflationary models.

PACS numbers:

I. INTRODUCTION

According to the Wilkinson Microwave Anisotropy Probe (WMAP) data [1] and other observations, it seems incontestable that the large scale structure of the universe stems from quantum fluctuations during the slow-roll inflation. Considering the accuracy of future observations, we expect higher order corrections to the simplest inflation model become important. Hence, it would be worth to study the corrections due to higher curvature terms to the slow-roll inflation. To the leading order, the corrections come from the Gauss-Bonnet and the axion-like parity violating coupling terms [2]. In fact, the same type of corrections can be derived from a particular superstring theory [3]. At present, however, we have not fully understood such corrections based on superstring theory. Hence, we have to resort to the phenomenological approach. We hope the understanding of higher curvature corrections to the inflationary scenario provides a clue to the study of superstring theory.

There have been many works concerning the Gauss-Bonnet and the parity violating corrections to the Einstein gravity. It is known that Gauss-Bonnet term can provide non-singular cosmological solutions [3, 4] and also provide alternative explanation for dark-energy [5, 6, 7, 8]. There are many works on scalar and tensor perturbations in Gauss-Bonnet cosmology [9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20]. The parity violating term has been considered in conjunction with the circular polarization of gravitational waves [21, 22, 23, 24, 25]. However, since no one has systematically investigated both terms simultaneously in the context of the slow-roll inflation, the combined effect on the slow-roll inflation is not so clear. Moreover, observational implications have not been discussed so far. Hence, a systematic study of higher curvature corrections to slow-roll inflationary scenario is desired.

In this paper, we study the slow-roll inflation with higher curvature corrections and derive the formula for the observables in terms of slow-roll parameters. It is stressed that the corrections could be relevant to observables in spite of the fact that its energetic contribution can be negligible. The reason is as follows. The Gauss-Bonnet term behaves as an effective potential term in the scalar field equation, though it does not contribute to the Friedmann equation in the slow-roll approximation. It reminds us of a hybrid model. That is why the effect could alter the tensor-to-scalar ratio. There is also an effect of the parity violating term on the gravitational waves as the leading effect. In fact, the circular polarization will be produced during the slow-roll inflation, which never occurs in the conventional inflationary scenario.

The organization of the paper is as follows: In section II, we define slow-roll parameters in the inflationary scenario with higher curvature corrections. In section III, we calculate the scalar spectral index. In section IV, we obtain the tensor spectral index and the tensor-to-scalar ratio. Here, we show the possibility of the enhancement of the tensor-to-scalar ratio due to the Gauss-Bonnet term. We also estimate the degree of the circular polarization of the primordial gravitational waves. In section V, the concrete examples are presented as an illustration. The final section is devoted to the conclusion.

*Electronic address: satoh@tap.scphys.kyoto-u.ac.jp

†Electronic address: jiro@tap.scphys.kyoto-u.ac.jp

II. SLOW-ROLL INFLATION

In this section, we will identify the slow-roll parameters. Since the parity violating term is not relevant to the background evolution, we have the corrections solely from the Gauss-Bonnet term. What we shall do is to characterize the correction in terms of the newly defined slow-roll parameters.

We consider the gravitational action with the Gauss-Bonnet and the parity violating terms

$$S = \frac{M_{\text{Pl}}^2}{2} \int d^4x \sqrt{-g} R - \int d^4x \sqrt{-g} \left[\frac{1}{2} \nabla^\mu \phi \nabla_\mu \phi + V(\phi) \right] - \frac{1}{16} \int d^4x \sqrt{-g} \xi(\phi) R_{\text{GB}}^2 + \frac{1}{16} \int d^4x \sqrt{-g} \omega(\phi) R \tilde{R} , \quad (1)$$

where $M_{\text{Pl}}^2 \equiv 1/8\pi G$ denotes the reduced Planck mass and $g_{\mu\nu}$ is the metric tensor. The Gauss-Bonnet term R_{GB} and the parity violating term $R\tilde{R}$ are defined by

$$R_{\text{GB}}^2 \equiv R^{\alpha\beta\gamma\delta} R_{\alpha\beta\gamma\delta} - 4R^{\alpha\beta} R_{\alpha\beta} + R^2 , \quad (2)$$

$$R\tilde{R} \equiv \frac{1}{2} \epsilon^{\alpha\beta\gamma\delta} R_{\alpha\beta\rho\sigma} R_{\gamma\delta}{}^{\rho\sigma} . \quad (3)$$

The inflaton field ϕ has the potential $V(\phi)$. We have introduced coupling functions $\xi(\phi)$ and $\omega(\phi)$. In principle, these functions should be calculated from the fundamental theory, such as superstring theory. Hence, it might be important to put constraints on these functions through observations.

Let us take the flat Friedmann-Robertson-Walker (FRW) metric

$$ds^2 = -dt^2 + a^2(t) \delta_{ij} dx^i dx^j , \quad (4)$$

where a is the scale factor. From the action (1), we obtain following background equations

$$3M_{\text{Pl}}^2 H^2 = \frac{1}{2} \dot{\phi}^2 + V + \frac{3}{2} H^3 \dot{\xi} \quad (5)$$

$$\ddot{\phi} + 3H\dot{\phi} + \frac{3}{2} H^2 (\dot{H} + H^2) \xi_{,\phi} + V_{,\phi} = 0 , \quad (6)$$

where a dot denotes the derivative with respect to the cosmic time, and H is the Hubble parameter defined by $H \equiv \dot{a}/a$. Note that coupling function ξ works as the effective potential for the inflaton ϕ , and only the Gauss-Bonnet coupling ξ appears in background equations because of parity symmetry of FRW metric. We will see the effect of the parity violating term appears only in tensor perturbations.

We consider the friction-dominated slow-roll inflation phase where the following inequalities hold:

$$\frac{\dot{\phi}^2}{M_{\text{Pl}}^2 H^2} \ll 1 , \quad \frac{H^3 \dot{\xi}}{M_{\text{Pl}}^2 H^2} = \frac{H \dot{\xi}}{M_{\text{Pl}}^2} \ll 1 , \quad \frac{\dot{H}}{H^2} \ll 1 , \quad \frac{\ddot{\phi}}{H \dot{\phi}} \ll 1 . \quad (7)$$

Under these circumstances, Eq.(5) reads

$$H^2 = \frac{V}{3M_{\text{Pl}}^2} . \quad (8)$$

This is nothing but the conventional slow-roll Friedman equation. While, the scalar field equation (6) becomes

$$\dot{\phi} = -\frac{V_{,\phi}}{3H} - \frac{1}{2} H^3 \xi_{,\phi} . \quad (9)$$

Here, the higher curvature correction plays a roll. If the last term can be neglected, we have the conventional slow-roll inflationary scenario. On the other hand, if the last term becomes important, the situation is different. Namely, the inflation is driven by the potential V , whereas the dynamics of the scalar field is determined by the Gauss-Bonnet term.

Now, we need to check the consistency conditions. The consistency condition for (7) reads

$$\frac{\dot{\phi}^2}{M_{\text{Pl}}^2 H^2} = \frac{M_{\text{Pl}}^2 V_{,\phi}^2}{V^2} + \frac{V_{,\phi} \xi_{,\phi}}{36 M_{\text{Pl}}^6} + \frac{V_{,\phi} \xi_{,\phi}}{3 M_{\text{Pl}}^2} \ll 1 \quad (10)$$

and

$$\frac{H^3 \dot{\xi}}{M_{Pl}^2 H^2} = \frac{H \xi_\phi \dot{\phi}}{M_{Pl}^2} = -\frac{\xi_{,\phi} V_{,\phi}}{3M_{Pl}^2} - \frac{H^4 \xi_\phi^2}{2M_{Pl}^2} \ll 1. \quad (11)$$

Third consistency condition for Eq.(7) leads to

$$\frac{\dot{H}}{H^2} = \frac{1}{2H} \frac{(H^2)'}{H^2} = \frac{\dot{\phi}}{2H} \frac{V_{,\phi}}{V} = -\frac{M_{Pl}^2}{2} \left(\frac{V_{,\phi}}{V} \right)^2 - \frac{1}{12} \frac{\xi_{,\phi} V_{,\phi}}{M_{Pl}^2} \ll 1. \quad (12)$$

Taking time derivative of Eq.(9), we get another consistency condition

$$\frac{\ddot{\phi}}{H\dot{\phi}} = \frac{M_{Pl}^2}{2} \frac{V_{,\phi}^2}{V^2} - M_{Pl}^2 \frac{V_{,\phi\phi}}{V} - \frac{V_{,\phi}\xi_{,\phi}}{4M_{Pl}^2} - \frac{V\xi_{,\phi\phi}}{6M_{Pl}^2} \ll 1. \quad (13)$$

For the slow-roll assumption to be valid, these terms must be sufficiently small. Hence, we define *five* slow-roll parameters,

$$\epsilon \equiv \frac{M_{Pl}^2}{2} \frac{V_{,\phi}^2}{V^2}, \quad \eta \equiv M_{Pl}^2 \frac{V_{,\phi\phi}}{V}, \quad \alpha \equiv \frac{V_{,\phi}\xi_{,\phi}}{4M_{Pl}^2}, \quad \beta \equiv \frac{V\xi_{,\phi\phi}}{6M_{Pl}^2}, \quad \gamma \equiv \frac{V^2 \xi_\phi^2}{18M_{Pl}^6} = \frac{4}{9} \frac{\alpha^2}{\epsilon}, \quad (14)$$

and impose the conditions $\epsilon, |\eta|, |\alpha|, |\beta|, \gamma \ll 1$. If these five conditions are satisfied, the inflaton field ϕ rolls down the potential slowly.

Note that, due to the Gauss-Bonnet term, the number of parameters and accompanying conditions increased from two for the conventional slow-roll inflation to five for this case. What we want to know is the effect of new parameters α, β, γ on the cosmological fluctuations.

For later convenience, we define ρ^2 and σ as

$$\rho^2 = \frac{1}{M_{Pl}^2} \frac{\dot{\phi}^2}{H^2} = 2\epsilon + \frac{4}{3}\alpha + \frac{\gamma}{2}, \quad \sigma = \frac{H\dot{\xi}}{M_{Pl}^2} = -\frac{4}{3}\alpha - \gamma. \quad (15)$$

It is also useful to notice the following relation

$$\frac{\dot{H}}{H^2} = -\frac{M_{Pl}^2}{2} \frac{V_{,\phi}^2}{V^2} - \frac{V_{,\phi}\xi_{,\phi}}{12M_{Pl}^2} = -\epsilon - \frac{\alpha}{3}. \quad (16)$$

From the next section, we study corrections to cosmological perturbations due to higher curvature terms.

III. SCALAR PERTURBATIONS

In this section, we study the higher curvature corrections to the scalar perturbations. It is convenient to use conformal time τ , defined as $a(t)d\tau = dt$, rather than the cosmic time t .

We consider the scalar perturbations A, B, ψ , and E defined by the perturbed metric

$$ds^2 = a^2(\tau) \left[-(1+2A)d\tau^2 + B_{|i}d\eta dx^i + (\delta_{ij} + 2\psi\delta_{ij} + 2E_{|ij})dx^i dx^j \right], \quad (17)$$

where the bar denotes the spatial derivative and the prime represents the derivative with respect to the conformal time. Here, we choose the gauge, $E = \delta\phi = 0$. In this gauge, modified Einstein equations for scalar quantities become

$$2(1-\sigma/2)\nabla^2\psi - 2\mathcal{H}(1-3\sigma/4)\nabla^2 B - 6\mathcal{H}(1-3\sigma/4)\psi' = 2a^2 V A / M_{Pl}^2 - 3\sigma\mathcal{H}^2 A \quad (18)$$

$$(1-3\sigma/4)\mathcal{H}A = -(1-\sigma/2)\psi' \quad (19)$$

$$(1-\sigma/2)A - \left(1 - \frac{\ddot{\xi}}{2M_{Pl}^2} \right) \psi + \frac{1}{a^2} [(1-\sigma/2)a^2 B]' = 0, \quad (20)$$

where $\mathcal{H}$ is conformal Hubble parameter, defined by $\mathcal{H} \equiv a'/a = aH$. By eliminating A and B from the above equations, we can get the single equation for ψ [10, 11]. It is also straightforward to obtain the action for ψ from the result.

Using Fourier expansion of ψ

$$\psi = \frac{1}{M_{\text{Pl}}} \int \frac{d^3k}{(2\pi)^3} \psi_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{x}} , \quad (21)$$

we can write down the action for $\psi_{\mathbf{k}}$

$$S = \frac{1}{2} \int d\tau \int \frac{d^3k}{(2\pi)^3} A_\psi^2 (|\psi'_{\mathbf{k}}|^2 - C_\psi^2 k^2 |\psi_{\mathbf{k}}|^2) , \quad (22)$$

where we have defined

$$A_\psi^2 \equiv 6a^2(1 - \sigma/2) \left[1 - (1 - \rho^2/6 - \sigma) \frac{1 - \sigma/2}{(1 - 3\sigma/4)^2} \right] , \quad (23)$$

$$C_\psi^2 \equiv \frac{2}{A_\psi^2} \left\{ \left[\frac{a(1 - \sigma/2)^2}{H(1 - 3\sigma/4)} \right]' - a^2 \left(1 - \frac{\ddot{\xi}}{2M_{\text{Pl}}^2} \right) \right\} . \quad (24)$$

Note that, due to the parity symmetry in scalar modes, there never appears the effect of the coupling function $\omega(\phi)$ in the action.

Defining a new variable $\Psi_{\mathbf{k}} \equiv A_\psi \psi_{\mathbf{k}}$, we obtain the canonical action

$$S = \frac{1}{2} \int d\tau \int \frac{d^3k}{(2\pi)^3} \left(|\Psi'_{\mathbf{k}}|^2 - C_\psi^2 k^2 |\Psi_{\mathbf{k}}|^2 + \frac{A''_\psi}{A_\psi} |\Psi_{\mathbf{k}}|^2 \right) . \quad (25)$$

This gives the Schrodinger type equation

$$\Psi''_{\mathbf{k}} + \left(C_\psi^2 k^2 - \frac{A''_\psi}{A_\psi} \right) \Psi_{\mathbf{k}} = 0 , \quad (26)$$

Notice that the term A''_ψ/A_ψ behaves as an effective potential, and C_ψ behaves as an effective sound speed.

We shall quantize the field $\Psi_{\mathbf{k}}$ by promoting it to the operator. We can use the mode expansion

$$\hat{\Psi}_{\mathbf{k}} = v_{\mathbf{k}}(\tau) \hat{a}_{\mathbf{k}} + v_{-\mathbf{k}}^*(\tau) \hat{a}_{-\mathbf{k}}^\dagger , \quad (27)$$

where $\hat{a}_{\mathbf{k}}$ and $\hat{a}_{-\mathbf{k}}^\dagger$ are annihilation and creation operators. The mode function $v_{\mathbf{k}}$ satisfies the same equation as $\Psi_{\mathbf{k}}$

$$v''_{\mathbf{k}} + \left(C_\psi^2 k^2 - \frac{A''_\psi}{A_\psi} \right) v_{\mathbf{k}} = 0 . \quad (28)$$

Under the slow-roll conditions, A_ψ can be approximated as

$$A_\psi^2 = 6a^2 \left[1 - (1 - \rho^2/6 - \sigma) \frac{1 - \sigma/2}{1 - 3\sigma/2} \right] = a^2 \rho^2 = a^2 (2\epsilon + 4\alpha/3 + \gamma/2) , \quad (29)$$

where we have kept terms to the leading order in slow-roll parameters. Similarly, C_ψ can be calculated as

$$C_\psi^2 = \frac{2}{A_\psi^2} \left[a \frac{d}{dt} \left(\frac{a}{H} \right) \frac{1 - \sigma}{1 - 3\sigma/4} - a^2 \left(1 - \frac{\ddot{\xi}}{2M_{\text{Pl}}^2} \right) \right] = \frac{2}{\rho^2} \left[\left(1 - \frac{\dot{H}}{H^2} \right) (1 - \sigma/4) - 1 + \frac{\ddot{\xi}}{2M_{\text{Pl}}^2} \right] . \quad (30)$$

The last term can be negligible as is seen from the following calculation

$$\ddot{\xi} = \xi_{,\phi\phi} \dot{\phi}^2 + \xi_{,\phi} \ddot{\phi} = M_{\text{Pl}}^2 H^2 \xi_{,\phi\phi} \rho^2 + H \dot{\xi} \frac{\ddot{\phi}}{H\dot{\phi}} = 2M_{\text{Pl}}^2 \beta \rho^2 + M_{\text{Pl}}^2 \sigma \frac{\ddot{\phi}}{H\dot{\phi}} .$$

Thus, we have

$$C_\psi^2 = \frac{2}{\rho^2} \left(-\frac{\dot{H}}{H^2} - \sigma/4 \right) = \frac{2(\epsilon + \alpha/3 + \alpha/3 + \gamma/4)}{2\epsilon + 4\alpha/3 + \gamma/2} = 1 + \mathcal{O}(\epsilon) , \quad (31)$$

where $\mathcal{O}(\epsilon)$ represents the first order corrections in slow-roll parameters. Now, let us calculate the effective potential. Using the relation

$$\dot{\rho} = -M_{\text{Pl}} \frac{V_{,\phi\phi}}{V} \dot{\phi} + M_{\text{Pl}} \frac{V_{,\phi}^2}{V^2} \dot{\phi} - \frac{1}{6M_{\text{Pl}}^3} V_{,\phi} \xi_{,\phi} \dot{\phi} - \frac{1}{6M_{\text{Pl}}^3} V \xi_{,\phi\phi} \dot{\phi} = \left(2\epsilon - \eta - \frac{2}{3}\alpha - \beta \right) H \rho , \quad (32)$$

we obtain

$$\frac{A'_\psi}{A_\psi} = \frac{a}{2} \frac{(A_\psi^2)'}{A_\psi^2} = a \left(H + \frac{\dot{\rho}}{\rho} \right) = \mathcal{H} \left(1 + 2\epsilon - \eta - \frac{2}{3}\alpha - \beta \right) .$$

From Eq.(16), we deduce

$$\left(\frac{1}{\mathcal{H}} \right)' = a \frac{d}{dt} \left(\frac{1}{aH} \right) = -\frac{\dot{H}}{H^2} - 1 = -1 - \epsilon - \frac{\alpha}{3} . \quad (33)$$

We integrate this equation to get $\mathcal{H}$ as

$$\mathcal{H} = \frac{1 + \epsilon + \alpha/3}{-\tau} , \quad (34)$$

where we assumed $\tau < 0$. Hence, we reached the following result

$$\frac{A'_\psi}{A_\psi} = \frac{(1 + \epsilon + \alpha/3)(1 + 2\epsilon - \eta - 2\alpha/3 - \beta)}{-\tau} = \frac{1 + 3\epsilon - \eta - \alpha/3 - \beta}{-\tau} . \quad (35)$$

Thus, we can get the effective potential

$$\frac{A''_\psi}{A_\psi} = \frac{d}{d\tau} \left(\frac{A'_\psi}{A_\psi} \right) + \left(\frac{A'_\psi}{A_\psi} \right)^2 = \frac{2 + 9\epsilon - 3\eta - \alpha - 3\beta}{\tau^2} = \frac{\nu_\psi^2 - 1/4}{\tau^2} , \quad (36)$$

where we have defined

$$\nu_\psi \equiv \frac{3}{2} + 3\epsilon - \eta - \alpha/3 - \beta . \quad (37)$$

Now, Eq.(28) reduces to

$$v_{\mathbf{k}}'' + \left(C_\psi^2 k^2 - \frac{\nu_\psi - 1/4}{\tau^2} \right) v_{\mathbf{k}} = 0 . \quad (38)$$

In the asymptotic past $\tau \rightarrow -\infty$, the effective potential vanishes. Hence, we can specify the Bunch-Davies vacuum by choosing the positive frequency mode function there as

$$v_{\mathbf{k}} = \frac{1}{\sqrt{2C_\psi k}} e^{-iC_\psi k\tau} . \quad (39)$$

It is easy to find the corresponding mode function

$$v_{\mathbf{k}}(\tau) = \frac{\sqrt{-\pi\tau}}{2} e^{i(\pi/4 + \pi\nu_\psi/2)} H_{\nu_\psi}^{(1)}(-C_\psi k\tau) , \quad (40)$$

where $H_{\nu_\psi}^{(1)}$ is the first kind of Hunkel function. Asymptotic form of $v_{\mathbf{k}}$ outside the horizon $-k\tau \ll 1$ reads

$$v_{\mathbf{k}}(\tau) = \sqrt{\frac{-\tau}{2(-C_\psi k\tau)^3}} e^{i(-\pi/4 + \pi\nu_\psi/2)} \frac{\Gamma(\nu_\psi)}{\Gamma(3/2)} \left(\frac{-C_\psi k\tau}{2} \right)^{3/2 - \nu_\psi} . \quad (41)$$

The power spectrum of scalar perturbation $\mathcal{P}_\psi$ can be defined as

$$\langle 0 | \hat{\psi}^\dagger \hat{\psi} | 0 \rangle = \int d(\log k) \frac{1}{2\pi^2} k^3 \frac{|v_{\mathbf{k}}|^2}{A_\psi^2} = \int d(\log k) \mathcal{P}_\psi(k) . \quad (42)$$

Thus, we obtain

$$\mathcal{P}_\psi(k) = \frac{1}{C_\psi^3} \frac{1}{4\pi^2} \frac{1}{A_\psi^2 \tau^2} \left(\frac{\Gamma(\nu_\psi)}{\Gamma(3/2)} \right)^2 \left(\frac{-C_\psi k \tau}{2} \right)^{3-2\nu_\psi}. \quad (43)$$

From Eqs.(29) and (34), we have

$$\frac{1}{A_\psi^2 \tau^2} = \frac{\mathcal{H}^2}{a^2(2\epsilon + 4\alpha/3 + \gamma/2)(1 + \epsilon + \alpha/3)^2} = \frac{H^2}{2\epsilon + 4\alpha/3 + \gamma/2}. \quad (44)$$

Using the above relation, we can simplify the power spectrum as

$$\mathcal{P}_\psi(k) = \frac{1}{2\epsilon + 4\alpha/3 + \gamma/2} \left(\frac{H}{2\pi} \frac{\Gamma(\nu_\psi)}{\Gamma(3/2)} \right)^2 \left(\frac{-k\tau}{2} \right)^{3-2\nu_\psi}. \quad (45)$$

Then, the scalar spectral index n_ψ is given by

$$n_\psi - 1 = 3 - 2\nu_\psi = -6\epsilon + 2\eta + \frac{2}{3}\alpha + 2\beta. \quad (46)$$

Of course, in the absence of the Gauss-Bonnet term, $\alpha = \beta = 0$, we recover the conventional formula.

IV. TENSOR PERTURBATIONS

In this section, we compute the power spectrum for tensor perturbations. In addition to the spectral index and the amplitude, we also calculate the circular polarization of gravitational waves.

The metric we consider is

$$ds^2 = a^2(\tau) [-d\tau^2 + (\delta_{ij} + h_{ij})dx^i dx^j], \quad (47)$$

where h_{ij} is a transverse and traceless tensor. Let us expand h_{ij} in plain waves and circular polarization tensor as

$$h_{ij} = \frac{\sqrt{2}}{M_{\text{Pl}}} \sum_{\pm} \int \frac{d^3k}{(2\pi)^3} \phi_{\mathbf{k}}^{\pm} e^{i\mathbf{k}\cdot\mathbf{x}} p_{\mathbf{k},ij}^{\pm}, \quad (48)$$

where $p_{\mathbf{k},ij}^{\pm}$ denotes polarization tensor for circular polarization, and $\pm$ denotes helicity of each mode. The action for $\phi_{\mathbf{k}}^{\pm}$ becomes,

$$S = \sum_{\pm} \frac{1}{2} \int d\tau \int \frac{d^3k}{(2\pi)^3} A_{\text{T}}^2 (|(\phi_{\mathbf{k}}^{\pm})'|^2 - C_{\text{T}}^2 k^2 |\phi_{\mathbf{k}}^{\pm}|^2), \quad (49)$$

where

$$A_{\text{T}}^2 \equiv a^2 \left(1 - \sigma/2 \mp \frac{1}{2M_{\text{Pl}}^2} \frac{k\omega'}{a^2} \right), \quad C_{\text{T}}^2 \equiv 1 + \frac{1}{2M_{\text{Pl}}^2} \frac{\mathcal{H}\xi'}{A_{\text{T}}^2} - \frac{1}{2M_{\text{Pl}}^2} \frac{a^2 \ddot{\xi}}{A_{\text{T}}^2}. \quad (50)$$

Note that there exists helicity dependence both in A_{T} and C_{T} , although we use notation A_{T} and C_{T} rather than $A_{\text{T}}^{\pm}$ and $C_{\text{T}}^{\pm}$, for simplicity. Apparently, the parity violating term gives rise to the left-right asymmetry, and hence the circular polarization of primordial gravitational waves. The observability of the circular polarization depends on the amplitude of the gravitational waves and the degree of the polarization. Since the parity violating term does not affect the background evolution and scalar perturbations, the coupling function ω cannot be constrained strongly. Here, we only require the absence of the ghost gravitons. From Eq.(50), we see a serious problem would appear if the parity violating term become dominant. In that case, $\pm$ sign appears as the overall factor of the action, and that means the appearance of ghost in the gravitational waves. In other words, we cannot trust our treatment of the gravitational waves in this regime. Hence, we impose that the term $|k\omega'/2a^2|$ must be lesser than M_{Pl}^2 at $k/a = M_c$,

$$1 > \frac{M_c}{2M_{\text{Pl}}^2} \left| \frac{\omega'}{a} \right| = \frac{M_c}{2M_{\text{Pl}}} \left| \frac{\dot{\omega}}{M_{\text{Pl}}} \right| \equiv |\Omega|, \quad (51)$$

where M_c is the physical wave-number corresponding to the cut-off scale, *e.g.* string scale, and we have defined

$$\Omega \equiv \frac{1}{2} \frac{M_c}{M_{\text{Pl}}} \frac{\dot{\omega}}{M_{\text{Pl}}} \simeq \text{const.} . \quad (52)$$

Here, since we are considering slow-roll inflation, we assumed $\dot{\omega} = \omega'/a \simeq \text{const.}$ in the period of interest. The amplitude of gravitational waves is determined at the horizon-crossing, $k/a \simeq H$. Thus, the effect of the parity violating term can be estimated as

$$\frac{1}{M_{\text{Pl}}^2} \left| \frac{k\omega'}{2a^2} \right| \simeq \frac{H}{2M_{\text{Pl}}^2} |\dot{\omega}| < \frac{H}{M_c} , \quad (53)$$

where we have used the constraint (51) in the last inequality. In other words, we can treat the parity violating effect as small correction always, since the Hubble parameter is smaller than the cut-off scale.

Defining new variables $\mu_{\mathbf{k}}^{\pm} \equiv A_{\text{T}} \phi_{\mathbf{k}}^{\pm}$, we obtain the canonical action

$$S = \sum_{\pm} \frac{1}{2} \int d\tau \int \frac{dk^3}{(2\pi)^3} \left(|(\mu_{\mathbf{k}}^{\pm})'|^2 - C_{\text{T}}^2 k^2 |\mu_{\mathbf{k}}^{\pm}|^2 + \frac{A_{\text{T}}''}{A_{\text{T}}} |\mu_{\mathbf{k}}^{\pm}|^2 \right) . \quad (54)$$

Taking the variation of the action with respect to $\mu_{\mathbf{k}}^{\pm}$, we obtain the equation for the gravitational waves

$$(\mu_{\mathbf{k}}^{\pm})'' + \left(C_{\text{T}}^2 k^2 - \frac{A_{\text{T}}''}{A_{\text{T}}} \right) \mu_{\mathbf{k}}^{\pm} = 0 . \quad (55)$$

Now, we quantize gravitational waves

$$\hat{h}_{ij} = \frac{\sqrt{2}}{M_{\text{Pl}}} \sum_{\pm} \int \frac{dk^3}{(2\pi)^3} \frac{\hat{\mu}_{\mathbf{k}}^{\pm}}{A_{\text{T}}} e^{i\mathbf{k} \cdot \mathbf{x}} p_{\mathbf{k},ij}^{\pm} , \quad (56)$$

by using the mode expansion

$$\hat{\mu}_{\mathbf{k}}^{\pm} = u_{\mathbf{k}}^{\pm}(\tau) \hat{a}_{\mathbf{k}}^{\pm} + u_{-\mathbf{k}}^{\pm*}(\tau) \hat{a}_{-\mathbf{k}}^{\pm\dagger} . \quad (57)$$

Let us represent A_{T} and C_{T} in terms of slow-roll parameters as

$$A_{\text{T}}^2 = a^2 \left(1 + \frac{2}{3} \alpha + \frac{\gamma}{2} \mp \frac{k\Omega}{M_c a} \right) \quad (58)$$

and

$$C_{\text{T}}^2 = 1 + \frac{1}{2M_{\text{Pl}}^2} \frac{\mathcal{H}\xi'}{A_{\text{T}}^2} = 1 + \frac{\sigma}{2(1 - \sigma/2 \mp k\Omega/M_c a)} = 1 - \frac{2}{3} \alpha - \frac{1}{2} \gamma . \quad (59)$$

Remember that $|\Omega| < 1$ from the constraint (51). Now, using the following relation

$$\frac{A_{\text{T}}'}{A_{\text{T}}} = \frac{1}{2} \frac{(A_{\text{T}}^2)'}{A_{\text{T}}^2} = \mathcal{H} \pm \frac{k\Omega}{2M_c} \frac{\mathcal{H}}{a} = \frac{1 + \epsilon + \alpha/3}{-\tau} \pm k \frac{H}{2M_c} \Omega , \quad (60)$$

we obtain the effective potential

$$\frac{A_{\text{T}}''}{A_{\text{T}}} = \frac{d}{d\tau} \left(\frac{A_{\text{T}}'}{A_{\text{T}}} \right) + \left(\frac{A_{\text{T}}'}{A_{\text{T}}} \right)^2 = \frac{2 + 3\epsilon + \alpha}{\tau^2} \pm \frac{k}{-\tau} \frac{H}{M_c} \Omega = \frac{\nu_{\text{T}}^2 - 1/4}{\tau^2} \pm \frac{k}{-\tau} \frac{H}{M_c} \Omega , \quad (61)$$

where we have defined

$$\nu_{\text{T}} \equiv \frac{3}{2} + \epsilon + \frac{\alpha}{3} . \quad (62)$$

Thus, the mode function $u_{\mathbf{k}}$ obeys

$$(u_{\mathbf{k}}^{\pm})'' + \left(C_{\text{T}}^2 k^2 - \frac{\nu_{\text{T}}^2 - 1/4}{\tau^2} \mp \frac{k}{-\tau} \frac{H}{M_c} \Omega \right) u_{\mathbf{k}}^{\pm} = 0 . \quad (63)$$

The Bunch-Davis vacuum is defined by the asymptotic form of the mode function at $\tau \rightarrow -\infty$

$$u_{\mathbf{k}}^{\pm} = \frac{1}{\sqrt{2C_{\text{T}}k}} e^{-iC_{\text{T}}k\tau} . \quad (64)$$

Using the confluent hypergeometric function U , the corresponding positive frequency mode function $u_{\mathbf{k}}^{\pm}$ is analytically given by

$$u_{\mathbf{k}}^{\pm}(\tau) = e^{-iC_{\text{T}}k\tau} (-2C_{\text{T}}k\tau)^{\nu_{\text{T}}} \sqrt{-\tau} e^{-i(\pi/4 + \pi\nu_{\text{T}}/2)} U\left(\frac{1}{2} + \nu_{\text{T}} \mp i\frac{1}{2C_{\text{T}}} \frac{H}{M_{\text{c}}} \Omega, 1 + 2\nu_{\text{T}}, 2iC_{\text{T}}k\tau\right) \\ \times \exp\left(\pm \frac{\pi}{4C_{\text{T}}} \frac{H}{M_{\text{c}}} \Omega\right) . \quad (65)$$

The asymptotic form of $u_{\mathbf{k}}^{\pm}$ outside the horizon $-C_{\text{T}}k\tau \ll 1$ becomes

$$u_{\mathbf{k}}^{\pm}(\tau) = \sqrt{\frac{-\tau}{2(-C_{\text{T}}k\tau)^3}} e^{i(-\pi/4 + \pi\nu_{\text{T}}/2)} \frac{\Gamma(\nu_{\text{T}})}{\Gamma(3/2)} \left(\frac{-C_{\text{T}}k\tau}{2}\right)^{3/2 - \nu_{\text{T}}} \exp\left(\pm \frac{\pi}{4C_{\text{T}}} \frac{H}{M_{\text{c}}} \Omega\right) . \quad (66)$$

Now, let us define the power spectrum $\mathcal{P}_{\text{T}}$ as

$$\langle 0 | \hat{h}_{ij} \hat{h}^{ij} | 0 \rangle = \sum_{\pm} \int d(\log k) \frac{2}{\pi^2} k^3 \frac{|u_{\mathbf{k}}^{\pm}|^2}{A_{\text{T}}^2} = \sum_{\pm} \int d(\log k) \mathcal{P}_{\text{T}}^{\pm} = \int d(\log k) \mathcal{P}_{\text{T}} . \quad (67)$$

Thus, we have the formula $\mathcal{P}_{\text{T}}^{\pm}$ for each polarization

$$\mathcal{P}_{\text{T}}^{\pm} = \frac{1}{C_{\text{T}}^3} \frac{1}{\pi^2} \frac{1}{A_{\text{T}}^2 \tau^2} \left(\frac{\Gamma(\nu_{\text{T}})}{\Gamma(3/2)}\right)^2 \left(\frac{-C_{\text{T}}k\tau}{2}\right)^{3-2\nu_{\text{T}}} \exp\left(\pm \frac{\pi}{2C_{\text{T}}} \frac{H}{M_{\text{c}}} \Omega\right) \\ = \frac{4}{1 + \epsilon - \alpha - \gamma} \left(\frac{H}{2\pi} \frac{\Gamma(\nu_{\text{T}})}{\Gamma(3/2)}\right)^2 \left(\frac{-C_{\text{T}}k\tau}{2}\right)^{3-2\nu_{\text{T}}} \left(1 \pm \frac{\pi}{2} \frac{H}{M_{\text{c}}} \Omega\right) \quad (68)$$

The total power spectrum $\mathcal{P}_{\text{T}} = \mathcal{P}_{\text{T}}^{+} + \mathcal{P}_{\text{T}}^{-}$ becomes

$$\mathcal{P}_{\text{T}} = \frac{8}{1 + \epsilon - \alpha - \gamma} \left(\frac{H}{2\pi} \frac{\Gamma(\nu_{\text{T}})}{\Gamma(3/2)}\right)^2 \left(\frac{-C_{\text{T}}k\tau}{2}\right)^{3-2\nu_{\text{T}}} . \quad (69)$$

It is easy to read off the tensor spectral index n_{T} as

$$n_{\text{T}} = 3 - 2\nu_{\text{T}} = -2\epsilon - \frac{2}{3}\alpha . \quad (70)$$

Again, this result reduces to the conventional slow-roll formula when $\alpha = 0$. The most impressive feature of the Gauss-Bonnet effects in the slow-roll inflation is shown in Eq.(70). In the usual case, because $\epsilon > 0$, n_{T} takes a negative value, namely, the tensor spectrum is red. However, if we incorporate the effect of the Gauss-Bonnet term, tensor spectrum could be blue, because α takes either positive or negative value. Hence, detection of the blue spectrum in the gravitational waves through the observation of B-mode polarization might indicate the existence of the Gauss-Bonnet term. In the case of the blue spectrum, $n_{\text{T}} > 0$, from Eq.(12), we see that the background evolution corresponds to superinflation $\dot{H} > 0$. In this case, inflaton field could climb up the potential slope, rather than roll down. This interesting possibility occurs due to the Gauss-Bonnet induced effective potential.

Since we have obtained both the scalar and tensor spectrum, we can deduce the tensor-to-scalar ratio r :

$$r \equiv \frac{\mathcal{P}_{\text{T}}}{\mathcal{P}_{\psi}} = 8 \frac{2\epsilon + 4\alpha/3 + \gamma/2}{1 + \epsilon - \alpha - \gamma} \left(\frac{-k\tau}{2}\right)^{2\nu_{\psi} - 2\nu_{\text{T}}} = \left(16\epsilon + \frac{32}{3}\alpha + 4\gamma\right) \left(\frac{k/a}{2H}\right)^{n_{\text{T}} - (n_{\psi} - 1)} . \quad (71)$$

Since we have the approximate relation $n_{\psi} - 1 = n_{\text{T}}$, the last factor must be 1. Hence, the tensor-to-scalar ratio r is given by

$$r \sim 16\epsilon + \frac{32}{3}\alpha + 4\gamma = 16\epsilon + \frac{32}{3}\alpha + \frac{16}{9} \frac{\alpha^2}{\epsilon} \geq \frac{32}{3} (\alpha + |\alpha|) . \quad (72)$$

For a negative α , we have the minimum, that is, $r = 0$. In the case $1 \geq \alpha \geq \epsilon$, the last term dominates and give rise to the sizable tensor-to-scalar ratio even if ϵ is extremely small.

Another interesting observable is the circular polarization Π , which is defined as difference between left and right helicity modes

$$\Pi \equiv \frac{\mathcal{P}_T^+ - \mathcal{P}_T^-}{\mathcal{P}_T^+ + \mathcal{P}_T^-} = \frac{\pi}{2} \frac{H}{M_c} \Omega. \quad (73)$$

One can see that the circular polarization is small under the constraint (51). However, it is important to quantify how small it is. Let us consider some extreme case, $H = 10^{-4} M_{\text{Pl}}$, $M_c = 10^{-3} M_{\text{Pl}}$, $\Omega = 0.1$, then, the ratio becomes $\Pi \simeq 0.015$. This means we have a hope to measure the sub-percent order of polarization due to the parity violating term [26, 27, 28]. It should be stressed that the effect of the Gauss-Bonnet term could enhance the amplitude of the primordial gravitational waves. That also enhances the detectability.

V. OBSERVATIONAL IMPLICATIONS

In the previous sections, we have derived general formula for observables in slow-roll inflation with higher curvature corrections. Here, we will discuss several models as an illustration.

Now, let us consider chaotic inflation models with higher curvature corrections. We take the functions as

$$V = \frac{1}{2} m^2 \phi^2, \quad \xi = \lambda e^{-\kappa \phi / M_{\text{Pl}}}, \quad (74)$$

where m is the mass of the inflaton and λ, κ are parameters of the coupling function ξ . Then, the slow-roll parameters can be calculated as

$$\epsilon = \eta = 2 \frac{M_{\text{Pl}}^2}{\phi^2}, \quad \alpha = -\frac{\kappa \lambda}{4} \frac{m^2 \phi}{M_{\text{Pl}}^3} e^{-\kappa \phi / M_{\text{Pl}}}, \quad \beta = \frac{\kappa^2 \lambda}{12} \frac{m^2 \phi^2}{M_{\text{Pl}}^4} e^{-\kappa \phi / M_{\text{Pl}}}, \quad \gamma = \frac{\kappa^2 \lambda^2}{72} \frac{m^4 \phi^4}{M_{\text{Pl}}^8} e^{-2\kappa \phi / M_{\text{Pl}}}. \quad (75)$$




FIG. 1: Expected spectral index n_ψ and tensor-to-scalar ratio r of the model (74) are shown. We have also plotted constraints from WMAP 5-year result, combined with BAO and Type I SN. The contours denote 68% and 95% confidence level. We used the out-put from Cosmological Parameters Plotter at LAMBDA [29]. Each line corresponds to the parameter (i) $m = 10^{-5} \times M_{\text{Pl}}$, $\lambda = 0$, $\kappa = 0$ (ordinary slow-roll), (ii) $m = 10^{-5} \times M_{\text{Pl}}$, $\lambda = 10^{10}$, $\kappa = 0.003$ and (iii) $m = 10^{-5} \times M_{\text{Pl}}$, $\lambda = 10^{10}$, $\kappa = 0.01$, from upper to lower, taking $\phi > 0$. Blue-colored lines denote the region that generate blue spectrum in tensor modes, and we plot some values of n_T .

We can evaluate the spectral index and the tensor-to-scalar ratio. The results are shown in FIG.1. There, we have plotted the scalar spectral index n_ψ and tensor-to-scalar ratio r of the model (74). We have also plotted constraints

from WMAP 5-year result, combined with baryon acoustic oscillations (BAO) and Type I supernova (SN). It turns out that our simple example explains WMAP results better than the conventional slow-roll inflation scenario. We see these observables are sensitive to higher curvature corrections. Hence, it implies that the higher curvature corrections are relevant to precision cosmology. Here, we notice that there exist parameter regions represented by blue lines which leads to the blue spectrum in tensor modes. We plotted some typical values of n_T in Fig.1. The e-folding numbers for the blue region cannot be estimated, because they correspond to superinflation. In that case, the model (74) must be modified so that the inflation ends.




FIG. 2: The tensor to the scalar ratio and the tensor spectral index are plotted. We have also shown the slow-roll parameters. Here we set parameters as $m = 10^{-5} \times M_{\text{Pl}}$, $\lambda = 10^{10}$ and $\kappa = 0.003$.

With the higher curvature corrections, there is no consistency relation as in the conventional inflation models. Therefore, in Fig.2, we have plotted the $r - n_T$ diagram. Since we have considered the negative α , the minimum of the tensor-to-scalar ratio is zero. Taking look at values for slow-roll parameters, we see the Gauss-Bonnet term dominates in the region $n_T \geq -0.01$.

We consider ϕ^4 model as another example. We parameterize the potential and the coupling functions as

$$V = \Lambda \phi^4, \quad \xi = \lambda e^{-\kappa \phi / M_{\text{Pl}}}, \quad (76)$$

where Λ is a coupling constant. Then, the slow-roll parameters read

$$\epsilon = 8 \frac{M_{\text{Pl}}^2}{\phi^2}, \quad \eta = 12 \frac{M_{\text{Pl}}^2}{\phi^2}, \quad \alpha = -\kappa \lambda \Lambda \frac{\phi^3}{M_{\text{Pl}}^3} e^{-\kappa \phi / M_{\text{Pl}}}, \quad \beta = \frac{\kappa^2 \lambda \Lambda}{6} \frac{\phi^4}{M_{\text{Pl}}^4} e^{-\kappa \phi / M_{\text{Pl}}}, \quad \gamma = \frac{\kappa^2 \lambda^2 \Lambda^2}{18} \frac{\phi^8}{M_{\text{Pl}}^8} e^{-2\kappa \phi / M_{\text{Pl}}}. \quad (77)$$

The scalar spectral index and the tensor-to-scalar ratio for some parameters are shown in FIG.3. We see that the ϕ^4 model with no higher curvature corrections cannot explain observations at all. However, if we take into account the higher curvature effects, the result changes and the ϕ^4 model seems to survive.

Let us give the last example where the tensor-to-scalar ratio r is enhanced due to the Gauss-Bonnet term. To highlight the feature, let us consider the flat potential model

$$V = V_0, \quad \xi = \lambda e^{-\kappa \phi / M_{\text{Pl}}}, \quad (78)$$

where V_0 is a constant. For this model, the slow-roll parameters are given by

$$\epsilon = \eta = \alpha = 0, \quad \beta = \frac{\kappa^2 \lambda}{6} \frac{V_0}{M_{\text{Pl}}^4} e^{-\kappa \phi / M_{\text{Pl}}}, \quad \gamma = \frac{\kappa^2 \lambda^2}{18} \frac{V_0^2}{M_{\text{Pl}}^8} e^{-2\kappa \phi / M_{\text{Pl}}}. \quad (79)$$

It is clear that r becomes zero and scalar spectrum becomes completely flat, if the higher curvature corrections are absent. Apparently, the spectral index $n_\psi = 1$ is not the favored value. Moreover, the detection of the primordial gravitational waves would observationally falsify this model. However, the higher curvature terms change the situation,




FIG. 3: Expected spectral index n_ψ and tensor-to-scalar ratio r of the model (76) are plotted. The constraints from WMAP 5-year data are also shown. Each line corresponds to the parameter (i) $\Lambda = 10^{-14}$, $\lambda = 0$, $\kappa = 0$ (ordinary slow-roll), (ii) $\Lambda = 10^{-14}$, $\lambda = 1.5 \times 10^{10}$, $\kappa = 0.1$ and (iii) $\Lambda = 10^{-14}$, $\lambda = 3.5 \times 10^{10}$, $\kappa = 0.1$, from upper to lower.

since β and γ are not zero! We show $r - n_\psi$ diagram, with correction terms, in FIG.4. Due to the higher curvature terms, the tensor-to-scalar ratio r becomes detectable and the scalar spectral index n_ψ is now a preferable value. This result clearly indicates the importance of higher curvature effects.




FIG. 4: The tensor-to-scalar ratio r and slow-roll parameters of the model (78) corresponding to scalar spectral index n_ψ are plotted. We set parameters as $V_0 = 10^{-10} \times M_{\text{Pl}}^4$, $\lambda = -10^{11}$ and $\kappa = 0.1$.

VI. CONCLUSION

We have studied the slow-roll inflationary scenario with the leading corrections, namely, the Gauss-Bonnet and the parity violating terms. We have obtained the higher curvature corrections to the scalar spectral index, the tensor

spectral index, and the tensor-to-scalar ratio. We found that the higher curvature corrections can not be negligible in the dynamics of the scalar field, although they are negligible in the Friedmann equation. It turned out that the tensor-to-scalar ratio could be enhanced because of the modified dynamics of the scalar field. We have found the tensor spectral index could be blue due to the Gauss-Bonnet term. We have also calculated the degree of circular polarization of gravitational waves generated during the slow-roll inflation. We discussed that the circular polarization can be observable due to the Gauss-Bonnet and parity violating terms. We also presented several examples to show observational implications of higher curvature corrections for chaotic inflationary models. We revealed that the observables are sensitive to the higher curvature corrections. Interestingly, there are some models which shows the slowly climbing up of the potential. Although it is an exotic possibility, the phenomena arises within the slow-roll approximation. The climbing up inflationary model deserves further investigation.

We can extend the present analysis in various ways. Although we have considered only the scalar field, we could have incorporated the vector field into the system [30, 31, 32], which leads to more general higher derivative corrections. However, the most important issue is to determine the coupling functions based on the fundamental physics such as the superstring theory. We leave these issues for future work.

Acknowledgments

We would like to thank Sugumi Kanno for the collaboration in the early stage of this project. J.S. is supported by the Japan-U.K. Research Cooperative Program, Grant-in-Aid for Scientific Research Fund of the Ministry of Education, Science and Culture of Japan No.18540262 and No.17340075.

-
- [1] E. Komatsu *et al.*, arXiv:0803.0547[astro-ph].
 - [2] S. Weinberg, arXiv:0804.4291[hep-th].
 - [3] I. Antoniadis, J. Rizos and K. Tamvakis, Nucl. Phys. B **415**, 497 (1994) [arXiv:hep-th/9305025].
 - [4] S. Kawai and J. Soda, Phys. Rev. D **59**, 063506 (1999) [arXiv:gr-qc/9807060].
 - [5] S. Nojiri, S. D. Odintsov and M. Sasaki, Phys. Rev. D **71**, 123509 (2005) [arXiv:hep-th/0504052].
 - [6] I. P. Neupane and B. M. N. Carter, JCAP **0606**, 004 (2006) [hep-th/0512262].
 - [7] I. P. Neupane, arXiv:0711.3234[hep-th].
 - [8] I. P. Neupane, Class. Quant. Grav. **23**, 7493 (2006) [hep-th/0602097].
 - [9] S. Kawai and J. Soda, arXiv:gr-qc/9906046.
 - [10] S. Kawai and J. Soda, Phys. Lett. B **460**, 41 (1999) [arXiv:gr-qc/9903017].
 - [11] J. c. Hwang and H. Noh, Phys. Rev. D **61**, 043511 (2000) [arXiv:astro-ph/9909480].
 - [12] C. Cartier, J. c. Hwang and E. J. Copeland, Phys. Rev. D **64**, 103504 (2001) [arXiv:astro-ph/0106197].
 - [13] S. Kawai, M. a. Sakagami and J. Soda, Phys. Lett. B **437**, 284 (1998) [arXiv:gr-qc/9802033].
 - [14] J. Soda, M. a. Sakagami and S. Kawai, arXiv:gr-qc/9807056.
 - [15] S. Kawai, M. a. Sakagami and J. Soda, arXiv:gr-qc/9901065.
 - [16] M. Gasperini, Phys. Rev. D **56**, 4815 (1997) [arXiv:gr-qc/9704045].
 - [17] C. Cartier, E. J. Copeland and M. Gasperini, Nucl. Phys. B **607**, 406 (2001) [arXiv:gr-qc/0101019].
 - [18] B. M. Leith and I. P. Neupane, JCAP **0705**, 019 (2007) [arXiv:hep-th/0702002].
 - [19] Z. K. Guo, N. Ohta and S. Tsujikawa, Phys. Rev. D **75**, 023520 (2007), [arXiv/hep-th:0610336].
 - [20] S. E. Vazquez, arXiv:0806.0603 [hep-th].
 - [21] A. Lue, L. M. Wang and M. Kamionkowski, Phys. Rev. Lett. **83**, 1506 (1999) [arXiv:astro-ph/9812088].
 - [22] K. Choi, J. c. Hwang and K. W. Hwang, Phys. Rev. D **61**, 084026 (2000) [arXiv:hep-ph/9907244].
 - [23] S. Alexander and J. Martin, Phys. Rev. D **71**, 063526 (2005) [arXiv:hep-th/0410230].
 - [24] D. H. Lyth, C. Quimbay and Y. Rodriguez, JHEP **0503**, 016 (2005) [arXiv:hep-th/0501153].
 - [25] M. Satoh, S. Kanno and J. Soda, Phys. Rev. D **77**, 023526 (2008), arXiv:0706.3585[astro-ph].
 - [26] S. Saito, K. Ichiki and A. Taruya, JCAP **0709**, 002 (2007), arXiv:0705.3701 [astro-ph].
 - [27] N. Seto, Phys. Rev. Lett. **97**, 151101 (2006) [arXiv:astro-ph/0609504].
 - [28] N. Seto and A. Taruya, Phys. Rev. Lett. **99**, 121101 (2007), arXiv:0707.0535 [astro-ph].
 - [29] <http://lambda.gsfc.nasa.gov/>
 - [30] K. Dimopoulos and M. Karcauskas, arXiv:0803.3041 [hep-th].
 - [31] S. Yokoyama and J. Soda, arXiv:0805.4265 [astro-ph].
 - [32] S. Kanno, M. Kimura, J. Soda and S. Yokoyama, arXiv:0806.2422 [hep-ph].